

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS.**

CAUTION

May be harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets and birds, and cover fish aquaria before space spraying or surface applications.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE.

Use only on dogs, cattle, goats, sheep, hogs, horses.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Do not expose to freezing temperatures. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility in accordance with PR Notice 83-3.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

CONTAINERS ONE GALLON AND SMALLER: Do not transport or store below 32 degree F. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

INDOOR APPLICATION

Do not use in food-handling establishments, restaurants or other areas where food is commercially prepared or processed. May be used in the non-food areas of USDA-inspected meat and poultry plants, as well as the non-food areas of other food-handling establishments. Non-Food/Feed Areas include (but are not limited to) garbage rooms, lavatories, floor drain (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, etc.

Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended. Retreat if reinfestation occurs but not more than once every 2 weeks.

SURFACE SPRAYING: To control **cockroaches, waterbugs, palmetto bugs, scorpions, millipedes, centipedes, sowbugs, pillbugs, ants, silverfish, firebrats, spiders, crickets, clover mites, cheese mites, granary weevils, confused flour beetles, rust red flour beetles, drugstore beetles, meal worms, grain mites and cadelles.** Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

To control **carpet beetles:** Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control **bedbugs:** Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Allow all treated articles to thoroughly dry before use. Repeat treatment as necessary.

To control **fleas and ticks** (adults and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed.

To control **clothes moths:** Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To kill **lice and louse eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying. Then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of the bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

Summit PERMACIDE® P-II

DIRECTIONS FOR USE Continued

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of **house flies, fruit flies, mosquitoes, skipper flies, wasps, hornets, bees, black flies, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths:** Direct spray at an upward angle, distribute it uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for 2 hours. Ventilate before reoccupying.

FOGGING: To kill exposed or accessible stage of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Ventilate before reoccupying.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from **horn flies, face flies, house flies, mosquitoes and gnats:** Apply a light mist sufficient to wet the surface of the hair. For **face flies** spray face and head, but do not spray into eyes.

To control **stable flies, horse flies and deer flies:** Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control **blood-sucking lice:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control **poultry lice:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control **bedbugs and mites** in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS: To control **fleas, ticks and lice:** Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Use protective gloves (mitts) to prevent contact with hands. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Certain medications can interact with pesticides. Consult a veterinarian before using on medicated animals.

NOTE: Do not use on animals under 12 weeks.

OUTDOOR APPLICATION

Do not allow spray treatment to drift on pasture land, cropland, poultry ranges or water supplies. Do not use on crops used for food, forage or pasture.

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill **Japanese beetles** hit by spray, as well as **whiteflies, aphids, exposed thrips, spiders, red mites and leafminers.**

May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas. May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geranium, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jew and zinnias.

Effective when used as a directive against **whiteflies, aphids, spider and red mites, exposed thrips, lace bugs, Japanese beetles, pavement ants, clover mites and army worms.** When spraying plants do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 degrees F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

TICKS AND ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where **ticks and mosquitoes** may rest. Apply at a rate of 2.5 pints per acre (0.9 fl. oz. per 1000square feet). Using truck-mounted conventional misting equipment such as rotomist models 51C, 91 or 100 HT traveling at 5 mph to treat a 300 foot swath, calibrate equipment to deliver 0.95 gallon per minute. Under no conditions should open water be treated with this product. Not to be used within 100 feet (30 meters) of lakes and streams.

Insecticides For The Food Industry

Summit PERMACIDE® P-II

For Use Outdoors and Non-Food Areas of Food/Feed Processing Establishments, Restaurants, Commercial Building: Warehouses, Theaters, Office Buildings, Schools, Motels, Hotels and Kennels

Authorized for Use in Federally Inspected Meat and Poultry Plants. Controls Fleas, Brown Dog Ticks and Lice On Premises and On Dogs.
Recommended For Commercial/Agricultural Use and Mosquito Abatement District Officials

ACTIVE INGREDIENTS:

*Premethrin.....	0.5%
*[3-phenoxyphenyl]methyl(+/-)cis:trans 3-(2,2-dichloroethenyl)2,2-dimethylcyclopropanecarboxylate.....	99.5%
Cis/trans ratio: min. 35%(+/-) cis and max. 65%(+/-) trans.	

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

- IF SWALLOWED:**
- * Call a poison control center or doctor immediately for treatment advice.
 - * Have a person sip a glass of water if able to swallow.
 - * Do not induce vomiting unless told to do so by the poison control center or doctor.
 - * Do not give anything by mouth to an unconscious person.
- IF ON SKIN OR CLOTHING:**
- * Take off contaminated clothing.
 - * Rinse skin immediately with plenty of water for 15-20 minutes.
- IF IN EYES:**
- * Call a poison control center or doctor for treatment advice.
 - * Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - * Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eye.
 - * Call a poison control center or doctor immediately for treatment advice.
- IF INHALED:**
- * Move person to fresh air.
 - * If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - * Call a poison control center or doctor for further treatment advice.
- Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.

EPA Registration No. 6218-64

A PRODUCT OF

EPA Establishment No. 6218-MD-2

BALTIMORE

MARYLAND



PERMACIDE® - Registered Trademark of Summit Chemical Co.

Summit **PERMACIDE® P-II**

For Use Outdoors and Non-Food Areas of Food/Feed Processing Establishments, Restaurants, Commercial Building: Warehouses, Theaters, Office Buildings, Schools, Motels, Hotels and Kennels

Authorized for Use in Federally Inspected Meat and Poultry Plants. Controls Fleas, Brown Dog Ticks and Lice On Premises and On Dogs.
Recommended For Commercial/Agricultural Use and Mosquito Abatement District Officials

ACTIVE INGREDIENTS:

*Premethrin..... 0.5%

INERT INGREDIENTS:

..... 99.5%

*(3-phenoxyphenyl)menthyl(+/-)cis/trans 3-(2,2-dichloroethyl)2,2-dimethylcyclopropanecarboxylate.

Cis/trans ratio: min. 35%(+/-) cis and max. 65%(+/-) trans.

KEEP OUT OF REACH OF CHILDREN

CAUTION**FIRST AID**

- IF SWALLOWED:**
- * Call a poison control center or doctor immediately for treatment advice.
 - * Have a person sip a glass of water if able to swallow.
 - * Do not induce vomiting unless told to do so by the poison control center or doctor.
 - * Do not give anything by mouth to an unconscious person.
- IF ON SKIN OR CLOTHING:**
- * Take off contaminated clothing.
 - * Rinse skin immediately with plenty of water for 15-20 minutes.
- IF IN EYES:**
- * Call a poison control center or doctor for treatment advice.
 - * Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - * Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eye.
 - * Call a poison control center or doctor immediately for treatment advice.
- IF INHALED:**
- * Move person to fresh air.
 - * If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - * Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.

EPA Registration No. 6218-64

A PRODUCT OF

EPA Establishment No. 6218-MD-2

BALTIMORE

MARYLAND



PERMACIDE® - Registered Trademark of Summit Chemical Co.



Peel Here

Net Contents: 1 Gallon

Peel Here

